

Fundamental solutions for a class of three-dimensional elliptic equations with singular coefficients

Anvar Hasanov and E.T. Karimov

*Institute of Mathematics and Information Technologies, Uzbek Academy of Sciences,
Durmon yuli str. 29, Tashkent 100125, Uzbekistan,
E-mail: anvarhasanov@yahoo.com, erkinjon@gmail.com*

Abstract

We consider an equation

$$L_{\alpha,\beta,\gamma}(u) \equiv u_{xx} + u_{yy} + u_{zz} + \frac{2\alpha}{x}u_x + \frac{2\beta}{y}u_y + \frac{2\gamma}{z}u_z = 0$$

in a domain $\mathbf{R}_3^+ \equiv \{(x, y, z) : x > 0, y > 0, z > 0\}$. Here α, β, γ are constants, moreover $0 < 2\alpha, 2\beta, 2\gamma < 1$. Main result of this paper is a construction of eight fundamental solutions for above-given equation in an explicit form. They are expressed by Lauricella's hypergeometric functions of three variables. Using expansion of Lauricella's hypergeometric function by products of Gauss's hypergeometric functions, it is proved that the found solutions have a singularity of the order $1/r$ at $r \rightarrow 0$. Furthermore, some properties of these solutions, which will be used at solving boundary-value problems for afore-mentioned equation are shown.

Keywords: Fundamental solutions, Elliptic differential equation with singular coefficients, Lauricella's hypergeometric function of three variables.

MSC 2000: *Primary* 35A08 *Secondary* 35J70

1 Introduction

Known that fundamental solutions (FSs) have an essential role at studying partial differential equations (PDEs). Formulation and solving of many local and non-local boundary-value problems (BVPs) are based on these solutions. Moreover, FSs appear as potentials, for instance, as a simple-layer and double-layer potentials in the theory of potentials.

Explicit form of FSs gives a possibility to study considered equation in detail. For example, in the works [3-5] by J.Barros-Neto and I.M.Gelfand, FSs for Tricomi operator, relative to an arbitrary point in the plane were explicitly calculated. Shown that found FSs clearly reflect the change of type of the Tricomi operator across the x -axis. We also mention J.Leray's work [12], where a general method, based upon the theory of analytic functions of several complex variables, for finding FSs for a class of hyperbolic linear differential operators with analytic coefficients was described. In particular, he showed how his method could be used to obtain, in the hyperbolic region, a FS for the Tricomi operator relative to a point $(0, b)$. Among other results on this direction we would like to note work by M.Itagaki [11], where three-dimensional high-order fundamental solutions for modified Helmholtz equation were found. Found solutions can be applied with the boundary particle method to some 2D inhomogeneous problems, for example, see [8].

Various modifications of the equation

$$L_{\alpha,\beta,\gamma}(u) \equiv u_{xx} + u_{yy} + u_{zz} + \frac{2\alpha}{x}u_x + \frac{2\beta}{y}u_y + \frac{2\gamma}{z}u_z = 0, \quad 0 < 2\alpha, 2\beta, 2\gamma < 1, \quad \alpha, \beta, \gamma = const, \quad (1)$$

in two-dimensional case were considered in many papers [1, 7, 13-18]. For example, in [7] the author is concerned with generalized bi-axially symmetric potentials (GBSP), i.e., with functions

$$u(x, y) = \sum_{n=0}^{\infty} a_n r^{2n} P_n^{(\alpha,\beta)}(\cos 2\theta),$$

where $x = r \cos \theta$, $y = r \sin \theta$, $\alpha, \beta > -1/2$, $P_n^{(\alpha, \beta)}(\cos 2\theta)$ are Jacobi polynomials. These functions occur as solutions of the equation

$$u_{xx} + u_{yy} + \frac{2\beta + 1}{x}u_x + \frac{2\alpha + 1}{y}u_y = 0.$$

Theorems on the convergence of above series and growth properties are obtained in terms of the coefficients a_n .

In [1], the elliptic or ultra hyperbolic equation

$$Lu = \sum_{i=1}^n \left(\frac{\partial^2 u}{\partial x_i^2} + \frac{\alpha_i}{x_i} \frac{\partial u}{\partial x_i} \right) \pm \sum_{i=1}^s \left(\frac{\partial^2 u}{\partial y_i^2} + \frac{\beta_i}{y_i} \frac{\partial u}{\partial y_i} \right) + \frac{\gamma u}{r^2} = 0, \quad (2)$$

is considered, where the constants $\alpha_1, \dots, \alpha_n, \beta_1, \dots, \beta_s, \gamma$ are real parameters, $r^2 = \sum_{i=1}^n x_i^2 \pm \sum_{i=1}^s y_i^2$ and establishes some properties of the operator L . An example is the Kelvin principle for the equation (2): If $u(x, y)$ is a solution of (2), then the function $\sigma = r^{-\varphi} u(\xi, \eta)$ is also a solution of the same equation, where $\xi_i = x_i/r^2$, $\eta_i = y_i/r^2$, $\varphi = n + s - 2 + \sum_{i=1}^n \alpha_i + \sum_{i=1}^s \beta_i$. For a class of iterated equations $L^p u = 0$, it was shown that if $Lu = 0$, then $L^p [r^{2j} u(x, y)] = L^p [r^{2j-\varphi} u(\xi, \eta)] = 0$ for $j = 0, \dots, p-1$; if, the function $u(x, y)$ is a homogeneous solution of degree λ of the equation $Lu = 0$, then $L^p [r^{2j-\varphi-2\lambda} u(x, y)] = L^p [r^{2(j+\lambda)} u(\xi, \eta)] = 0$ for $j = 0, 1, \dots, p-1$.

FSs for elliptic equations with singular coefficients are directly connected with hypergeometric functions (HFs). Therefore, basic properties such as decomposition formulas, integral representations, formulas of analytical continuation, formulas of differentiation for HFs are necessary for studying FSs.

Since afore-mentioned properties of HFs of Gauss, Appell, Kummer were well-known [2], results on investigations of elliptic equations with one or two singular coefficients were successful. Up to publication results on Lauricella HFs by A.Hasanov and H.M.Srivastava [9, 10] there was no possibility to investigate FSs for the equation (1), since their consist Lauricella HFs of three variables.

In the present paper we construct eight fundamental solutions for the equation (1) in an explicit form. Using decomposition formulas for Lauricella HFs of three variables by simple Gauss HFs, we prove that found FSs have a singularity of order $1/r$ at $r \rightarrow 0$.

The plan of this paper is as follows. In Section 2 we briefly give some preliminary information, which will be used further. Also some constructive formulas for the operator $L_{\alpha, \beta, \gamma}$ are given. In section 3 we describe the method of finding FSs for considered equation and show what order of singularity will have found solutions.

2 Preliminaries

Below we give some formulas for Euler gamma-function, Gauss HF, Lauricella HF of three variables, which will be used in next section.

Known that Euler gamma-function $\Gamma(a)$ has properties ([6], pp.17-19, (2), (10), (15))

$$\Gamma(a + m) = \Gamma(a)(a)_m; \quad \Gamma(a + 1/2) = \frac{\sqrt{\pi}\Gamma(2a)}{2^{2a-1}\Gamma(a)}, \quad \Gamma(1/2) = \sqrt{\pi}. \quad (3)$$

Here $(a)_m$ is a Pochhammer symbol, for which an equality $(a)_{m+n} = (a)_m(a+m)_n$ is true ([6], p.67, (5)).

A function ${}_2F_1(a, b; c; x) = \sum_{n=0}^{\infty} \frac{(a)_n(b)_n}{(c)_n n!} x^n$ is known as Gauss HF and an equality

$${}_2F_1(a, b; c; 1) = \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-b)\Gamma(c-a)}, \quad c \neq 0, -1, -2, \dots, \quad \operatorname{Re}(c-a-b) > 0 \quad (4)$$

is hold ([6], p.73, (14)). Moreover, the following autotransformer formula ([6], p.76, (22))

$${}_2F_1(a, b; c; x) = (1-x)^{-b} {}_2F_1(c-a, b; c; \frac{x}{x-1}) \quad (5)$$

is valid.

The following system called as the system of hypergeometric equations of Lauricella ([2], p. 117):

$$\begin{cases} \xi(1-\xi)\omega_{\xi\xi} - \xi\eta\omega_{\xi\eta} - \xi\zeta\omega_{\xi\zeta} + [2\alpha - (\alpha + \beta + \gamma + \frac{1}{2} + \alpha + 1)\xi]\omega_{\xi} - \alpha\eta\omega_{\eta} \\ \quad - \alpha\zeta\omega_{\zeta} - (\alpha + \beta + \gamma + \frac{1}{2})\alpha\omega = 0, \\ \eta(1-\eta)\omega_{\eta\eta} - \xi\eta\omega_{\xi\eta} - \eta\zeta\omega_{\eta\zeta} + [2\beta - (\alpha + \beta + \gamma + \frac{1}{2} + \beta + 1)\eta]\omega_{\eta} - \beta\xi\omega_{\xi} \\ \quad - \beta\zeta\omega_{\zeta} - (\alpha + \beta + \gamma + \frac{1}{2})\beta\omega = 0, \\ \zeta(1-\zeta)\omega_{\zeta\zeta} - \xi\zeta\omega_{\xi\zeta} - \eta\zeta\omega_{\eta\zeta} + [2\gamma - (\alpha + \beta + \gamma + \frac{1}{2} + \gamma + 1)\zeta]\omega_{\zeta} - \gamma\xi\omega_{\xi} \\ \quad - \gamma\eta\omega_{\eta} - (\alpha + \beta + \gamma + \frac{1}{2})\gamma\omega = 0. \end{cases} \quad (6)$$

The system (6) has the following solutions ([2], p. 118):

$$\omega_1(\xi, \eta, \zeta) = F_A^{(3)}\left(\alpha + \beta + \gamma + \frac{1}{2}; \alpha, \beta, \gamma; 2\alpha, 2\beta, 2\gamma; \xi, \eta, \zeta\right), \quad (7)$$

$$\omega_2(\xi, \eta, \zeta) = \xi^{1-2\alpha} F_A^{(3)}\left(-\alpha + \beta + \gamma + \frac{3}{2}; 1 - \alpha, \beta, \gamma; 2 - 2\alpha, 2\beta, 2\gamma; \xi, \eta, \zeta\right), \quad (8)$$

$$\omega_3(\xi, \eta, \zeta) = \eta^{1-2\beta} F_A^{(3)}\left(\alpha - \beta + \gamma + \frac{3}{2}; \alpha, 1 - \beta, \gamma; 2\alpha, 2 - 2\beta, 2\gamma; \xi, \eta, \zeta\right), \quad (9)$$

$$\omega_4(\xi, \eta, \zeta) = \zeta^{1-2\gamma} F_A^{(3)}\left(\alpha + \beta - \gamma + \frac{3}{2}; \alpha, \beta, 1 - \gamma; 2\alpha, 2\beta, 2 - 2\gamma; \xi, \eta, \zeta\right), \quad (10)$$

$$\omega_5(\xi, \eta, \zeta) = \xi^{1-2\alpha} \eta^{1-2\beta} F_A^{(3)}\left(-\alpha - \beta + \gamma + \frac{5}{2}; 1 - \alpha, 1 - \beta, \gamma; 2 - 2\alpha, 2 - 2\beta, 2\gamma; \xi, \eta, \zeta\right), \quad (11)$$

$$\omega_6(\xi, \eta, \zeta) = \xi^{1-2\alpha} \zeta^{1-2\gamma} F_A^{(3)}\left(-\alpha + \beta - \gamma + \frac{5}{2}; 1 - \alpha, \beta, 1 - \gamma; 2 - 2\alpha, 2\beta, 2 - 2\gamma; \xi, \eta, \zeta\right), \quad (12)$$

$$\omega_7(\xi, \eta, \zeta) = \eta^{1-2\beta} \zeta^{1-2\gamma} F_A^{(3)}\left(\alpha - \beta - \gamma + \frac{5}{2}; \alpha, 1 - \beta, 1 - \gamma; 2\alpha, 2 - 2\beta, 2 - 2\gamma; \xi, \eta, \zeta\right), \quad (13)$$

$$\begin{aligned} \omega_8(\xi, \eta, \zeta) \\ = \xi^{1-2\alpha} \eta^{1-2\beta} \zeta^{1-2\gamma} F_A^{(3)}\left(-\alpha - \beta - \gamma + \frac{7}{2}; 1 - \alpha, 1 - \beta, 1 - \gamma; 2 - 2\alpha, 2 - 2\beta, 2 - 2\gamma; \xi, \eta, \zeta\right), \end{aligned} \quad (14)$$

where ([2], pp. 114 - 115 (1), (5))

$$F_A^{(3)}(a; b_1, b_2, b_3; c_1, c_2, c_3; x, y, z) = \sum_{i,j,k=0}^{\infty} \frac{(a)_{i+j+k} (b_1)_i (b_2)_j (b_3)_k}{(c_1)_i (c_2)_j (c_3)_k i! j! k!} x^i y^j z^k, \quad (|x| + |y| + |z| < 1),$$

$$\begin{aligned} F_A^{(3)}(a; b_1, b_2, b_3; c_1, c_2, c_3; x, y, z) &= \frac{\Gamma(c_1)\Gamma(c_2)\Gamma(c_3)}{\Gamma(b_1)\Gamma(b_2)\Gamma(b_3)\Gamma(c_1 - b_1)\Gamma(c_2 - b_2)\Gamma(c_3 - b_3)} \\ &\times \int_0^1 \int_0^1 \int_0^1 t_1^{b_1-1} t_2^{b_2-1} t_3^{b_3-1} (1-t_1)^{c_1-b_1-1} (1-t_2)^{c_2-b_2-1} (1-t_3)^{c_3-b_3-1} (1-x t_1 - y t_2 - z t_3)^{-a} dt_1 dt_2 dt_3, \\ &\text{Re } c_i > \text{Re } b_i > 0, \quad i = 1, 2, 3. \end{aligned}$$

The following decomposition formula ([9], p. 117, (14))

$$\begin{aligned} F_A^{(3)}(a; b_1, b_2, b_3; c_1, c_2, c_3; x, y, z) &= \sum_{l,m,n=0}^{\infty} \frac{(a)_{l+m+n} (b_1)_{l+m} (b_2)_{l+n} (b_3)_{m+n}}{(c_1)_{l+m} (c_2)_{l+n} (c_3)_{m+n} l! m! n!} x^{l+m} y^{l+n} z^{m+n} \\ &\times {}_2F_1(a + l + m, b_1 + l + m; c_1 + l + m; x) {}_2F_1(a + l + m + n, b_2 + l + n; c_2 + l + n; y) \\ &\times {}_2F_1(a + l + m + n, b_3 + m + n; c_3 + m + n; z) \end{aligned} \quad (15)$$

and formula of differentiation ([2], p.19, (20))

$$\frac{\partial^{i+j+k} F_A^{(3)}}{\partial x^i \partial y^j \partial z^k} = \frac{(a)_{i+j+k} (b_1)_i (b_2)_j (b_3)_k}{(c_1)_i (c_2)_j (c_3)_k} F_A^{(3)}(a + i + j + k, b_1 + i, b_2 + j, b_3 + k; c_1 + i, c_2 + j, c_3 + k; x, y, z) \quad (16)$$

are valid.

One of the reasons why we consider the equation (1) is an existing of constructive formulas for the operator $L_{\alpha,\beta,\gamma}$, which give a possibility to investigate the operator at various values of parameters α, β, γ .

Remark 1. *The following constructive formulas*

$$\begin{aligned} L_{\alpha,\beta,\gamma}(x^{1-2\alpha}u) &\equiv x^{1-2\alpha}L_{1-\alpha,\beta,\gamma}(u), \\ L_{\alpha,\beta,\gamma}(y^{1-2\beta}u) &\equiv y^{1-2\beta}L_{\alpha,1-\beta,\gamma}(u), \\ L_{\alpha,\beta,\gamma}(z^{1-2\gamma}u) &\equiv z^{1-2\gamma}L_{\alpha,\beta,1-\gamma}(u), \end{aligned} \quad (17)$$

$$\begin{aligned} L_{\alpha,\beta,\gamma}(x^{1-2\alpha}y^{1-2\beta}u) &\equiv x^{1-2\alpha}y^{1-2\beta}L_{1-\alpha,1-\beta,\gamma}(u), \\ L_{\alpha,\beta,\gamma}(x^{1-2\alpha}z^{1-2\gamma}u) &\equiv x^{1-2\alpha}z^{1-2\gamma}L_{1-\alpha,\beta,1-\gamma}(u), \\ L_{\alpha,\beta,\gamma}(y^{1-2\beta}z^{1-2\gamma}u) &\equiv y^{1-2\beta}z^{1-2\gamma}L_{\alpha,1-\beta,1-\gamma}(u), \end{aligned} \quad (18)$$

$$L_{\alpha,\beta,\gamma}(x^{1-2\alpha}y^{1-2\beta}z^{1-2\gamma}u) \equiv x^{1-2\alpha}y^{1-2\beta}z^{1-2\gamma}L_{1-\alpha,1-\beta,1-\gamma}(u) \quad (19)$$

are true.

Let us prove (19). Note, equalities (17), (18) can be proved similarly. First we find $L_{\alpha,\beta,\gamma}(x^{1-2\alpha}y^{1-2\beta}z^{1-2\gamma}u)$

$$\begin{aligned} &= (x^{1-2\alpha}y^{1-2\beta}z^{1-2\gamma}u)_{xx} + (x^{1-2\alpha}y^{1-2\beta}z^{1-2\gamma}u)_{yy} + (x^{1-2\alpha}y^{1-2\beta}z^{1-2\gamma}u)_{zz} \\ &+ \frac{2\alpha}{x}(x^{1-2\alpha}y^{1-2\beta}z^{1-2\gamma}u)_x + \frac{2\beta}{y}(x^{1-2\alpha}y^{1-2\beta}z^{1-2\gamma}u)_y + \frac{2\gamma}{z}(x^{1-2\alpha}y^{1-2\beta}z^{1-2\gamma}u)_z. \end{aligned}$$

Since

$$\begin{aligned} &(x^{1-2\alpha}y^{1-2\beta}z^{1-2\gamma}u)_{xx} \\ &= x^{1-2\alpha}y^{1-2\beta}z^{1-2\gamma}u_{xx} + 2(1-2\alpha)x^{-2\alpha}y^{1-2\beta}z^{1-2\gamma}u_x - 2\alpha(1-2\alpha)x^{-2\alpha-1}y^{1-2\beta}z^{1-2\gamma}u, \\ &(x^{1-2\alpha}y^{1-2\beta}z^{1-2\gamma}u)_{yy} \\ &= x^{1-2\alpha}y^{1-2\beta}z^{1-2\gamma}u_{yy} + 2(1-2\beta)x^{1-2\alpha}y^{-2\beta}z^{1-2\gamma}u_y - 2\beta(1-2\beta)x^{1-2\alpha}y^{-2\beta-1}z^{1-2\gamma}u, \\ &(x^{1-2\alpha}y^{1-2\beta}z^{1-2\gamma}u)_{zz} \\ &= x^{1-2\alpha}y^{1-2\beta}z^{1-2\gamma}u_{zz} + 2(1-2\gamma)x^{1-2\alpha}y^{1-2\beta}z^{-2\gamma}u_z - 2\gamma(1-2\gamma)x^{1-2\alpha}y^{1-2\beta}z^{-2\gamma-1}u, \\ &\frac{2\alpha}{x}(x^{1-2\alpha}y^{1-2\beta}z^{1-2\gamma}u)_x = 2\alpha x^{-2\alpha}y^{1-2\beta}z^{1-2\gamma}u_x + 2\alpha(1-2\alpha)x^{-2\alpha-1}y^{1-2\beta}z^{1-2\gamma}u, \\ &\frac{2\beta}{y}(x^{1-2\alpha}y^{1-2\beta}z^{1-2\gamma}u)_y = 2\beta x^{1-2\alpha}y^{-2\beta}z^{1-2\gamma}u_y + 2\beta(1-2\beta)x^{1-2\alpha}y^{-2\beta-1}z^{1-2\gamma}u, \\ &\frac{2\gamma}{z}(x^{1-2\alpha}y^{1-2\beta}z^{1-2\gamma}u)_z = 2\gamma x^{1-2\alpha}y^{1-2\beta}z^{-2\gamma}u_z + 2\gamma(1-2\gamma)x^{1-2\alpha}y^{1-2\beta}z^{-2\gamma-1}u, \end{aligned}$$

after some simplifications we get

$$\begin{aligned} &L_{\alpha,\beta,\gamma}(x^{1-2\alpha}y^{1-2\beta}z^{1-2\gamma}u) \\ &= x^{1-2\alpha}y^{1-2\beta}z^{1-2\gamma} \left(u_{xx} + u_{yy} + u_{zz} + \frac{2(1-\alpha)}{x}u_x + \frac{2(1-\beta)}{y}u_y + \frac{2(1-\gamma)}{z}u_z \right) \\ &= x^{1-2\alpha}y^{1-2\beta}z^{1-2\gamma}L_{1-\alpha,1-\beta,1-\gamma}(u). \end{aligned}$$

3 Finding fundamental solutions for the equation (1)

Solutions of the equation (1) we shall search as follows

$$u = P(r) \omega(\xi, \eta, \zeta), \quad (20)$$

where

$$\begin{aligned} r^2 &= (x - x_0)^2 + (y - y_0)^2 + (z - z_0)^2, \\ r_1^2 &= (x + x_0)^2 + (y - y_0)^2 + (z - z_0)^2, \\ r_2^2 &= (x - x_0)^2 + (y + y_0)^2 + (z - z_0)^2, \\ r_3^2 &= (x - x_0)^2 + (y - y_0)^2 + (z + z_0)^2, \end{aligned} \quad (21)$$

$$\xi = \frac{r^2 - r_1^2}{r^2}, \eta = \frac{r^2 - r_2^2}{r^2}, \zeta = \frac{r^2 - r_3^2}{r^2}, \quad P(r) = (r^2)^{-\alpha-\beta-\gamma-\frac{1}{2}}. \quad (22)$$

Here $\forall (x, y, z) \in \mathbf{R}_3^+$, but $(x_0, y_0, z_0) \in \mathbf{R}_3^+$ is any fixed point. Substituting (20) into (1) we get

$$A_1 \omega_{\xi\xi} + A_2 \omega_{\eta\eta} + A_3 \omega_{\zeta\zeta} + B_1 \omega_{\xi\eta} + B_2 \omega_{\xi\zeta} + B_3 \omega_{\eta\zeta} + C_1 \omega_\xi + C_2 \omega_\eta + C_3 \omega_\zeta + D \omega = 0, \quad (23)$$

where

$$\begin{aligned} A_1 &= P[\xi_x^2 + \xi_y^2 + \xi_z^2], \quad A_2 = P[\eta_x^2 + \eta_y^2 + \eta_z^2], \quad A_3 = P[\zeta_x^2 + \zeta_y^2 + \zeta_z^2], \\ B_1 &= 2P[\xi_x \eta_x + \xi_y \eta_y + \xi_z \eta_z], \quad B_2 = 2P[\xi_x \zeta_x + \xi_y \zeta_y + \xi_z \zeta_z], \quad B_3 = 2P[\eta_x \zeta_x + \eta_y \zeta_y + \eta_z \zeta_z], \\ C_1 &= P(\xi_{xx} + \xi_{yy} + \xi_{zz}) + 2(P_x \xi_x + P_y \xi_y + P_z \xi_z) + P\left(\xi_x \frac{2\alpha}{x} + \xi_y \frac{2\beta}{y} + \xi_z \frac{2\gamma}{z}\right), \\ C_2 &= P(\eta_{xx} + \eta_{yy} + \eta_{zz}) + 2(P_x \eta_x + P_y \eta_y + P_z \eta_z) + P\left(\eta_x \frac{2\alpha}{x} + \eta_y \frac{2\beta}{y} + \eta_z \frac{2\gamma}{z}\right), \\ C_3 &= P(\zeta_{xx} + \zeta_{yy} + \zeta_{zz}) + 2(P_x \zeta_x + P_y \zeta_y + P_z \zeta_z) + P\left(\zeta_x \frac{2\alpha}{x} + \zeta_y \frac{2\beta}{y} + \zeta_z \frac{2\gamma}{z}\right), \\ D &= P_{xx} + P_{yy} + P_{zz} + P_x \frac{2\alpha}{x} + P_y \frac{2\beta}{y} + P_z \frac{2\gamma}{z}. \end{aligned}$$

After several evaluations we find

$$A_1 = -4P(r^2)^{-1} x^{-1} x_0 \xi (1 - \xi), \quad A_2 = -4P(r^2)^{-1} y^{-1} y_0 \eta (1 - \eta), \quad (24)$$

$$A_3 = -4P(r^2)^{-1} z^{-1} z_0 \zeta (1 - \zeta), \quad B_1 = 4P(r^2)^{-1} x^{-1} x_0 \xi \eta + 4P(r^2)^{-1} y^{-1} y_0 \xi \eta, \quad (25)$$

$$B_2 = 4P(r^2)^{-1} x^{-1} x_0 \xi \zeta + 4P(r^2)^{-1} z^{-1} z_0 \xi \zeta, \quad B_3 = 4P(r^2)^{-1} y^{-1} y_0 \eta \zeta + 4P(r^2)^{-1} z^{-1} z_0 \eta \zeta, \quad (26)$$

$$C_1 = -4P(r^2)^{-1} x^{-1} x_0 [2\alpha - (\alpha + \beta + \gamma + \frac{1}{2} + \alpha + 1) \xi] + 4P(r^2)^{-1} y^{-1} y_0 \beta \xi + 4P(r^2)^{-1} z^{-1} z_0 \gamma \xi, \quad (27)$$

$$C_2 = -4P(r^2)^{-1} y^{-1} y_0 [2\beta - (\alpha + \beta + \gamma + \frac{1}{2} + \beta + 1) \eta] + 4P(r^2)^{-1} x^{-1} x_0 \alpha \eta + 4P(r^2)^{-1} z^{-1} z_0 \gamma \eta, \quad (28)$$

$$C_3 = -4P(r^2)^{-1} z^{-1} z_0 [2\gamma - (\alpha + \beta + \gamma + \frac{1}{2} + \gamma + 1) \zeta] + 4P(r^2)^{-1} x^{-1} x_0 \alpha \zeta + 4P(r^2)^{-1} y^{-1} y_0 \beta \zeta, \quad (29)$$

$$D = 4P(r^2)^{-1} x^{-1} x_0 (\alpha + \beta + \gamma + \frac{1}{2}) \alpha + 4P(r^2)^{-1} y^{-1} y_0 (\alpha + \beta + \gamma + \frac{1}{2}) \beta + 4P(r^2)^{-1} z^{-1} z_0 (\alpha + \beta + \gamma + \frac{1}{2}) \gamma. \quad (30)$$

Substituting equalities (24) - (30) into (23) we obtain the system of hypergeometric equations of Lauricella (6), which has solutions (7) - (14).

Considering (7) - (14), from (20) we obtain eight FSs of the equation (1)

$$q_1(x, y, z; x_0, y_0, z_0) = k_1 (r^2)^{-\alpha-\beta-\gamma-\frac{1}{2}} F_A^{(3)} \left(\alpha + \beta + \gamma + \frac{1}{2}; \alpha, \beta, \gamma; 2\alpha, 2\beta, 2\gamma; \xi, \eta, \zeta \right), \quad (31)$$

$$q_2(x, y, z; x_0, y_0, z_0) = k_2 (r^2)^{\alpha-\beta-\gamma-\frac{3}{2}} x^{1-2\alpha} x_0^{1-2\alpha} F_A^{(3)} \left(-\alpha + \beta + \gamma + \frac{3}{2}; 1 - \alpha, \beta, \gamma; 2 - 2\alpha, 2\beta, 2\gamma; \xi, \eta, \zeta \right), \quad (32)$$

$$q_3(x, y, z; x_0, y_0, z_0) = k_3 (r^2)^{-\alpha+\beta-\gamma-\frac{3}{2}} y^{1-2\beta} y_0^{1-2\beta} F_A^{(3)} \left(\alpha - \beta + \gamma + \frac{3}{2}; \alpha, 1 - \beta, \gamma; 2\alpha, 2 - 2\beta, 2\gamma; \xi, \eta, \zeta \right), \quad (33)$$

$$q_4(x, y, z; x_0, y_0, z_0) = k_4 (r^2)^{-\alpha-\beta+\gamma-\frac{3}{2}} z^{1-2\gamma} z_0^{1-2\gamma} F_A^{(3)} \left(\alpha + \beta - \gamma + \frac{3}{2}; \alpha, \beta, 1 - \gamma; 2\alpha, 2\beta, 2 - 2\gamma; \xi, \eta, \zeta \right), \quad (34)$$

$$q_5(x, y, z; x_0, y_0, z_0) = k_5 (r^2)^{\alpha+\beta-\gamma-\frac{5}{2}} \times x^{1-2\alpha} y^{1-2\beta} x_0^{1-2\alpha} y_0^{1-2\beta} F_A^{(3)} \left(-\alpha - \beta + \gamma + \frac{5}{2}; 1 - \alpha, 1 - \beta, \gamma; 2 - 2\alpha, 2 - 2\beta, 2\gamma; \xi, \eta, \zeta \right), \quad (35)$$

$$q_6(x, y, z; x_0, y_0, z_0) = k_6 (r^2)^{\alpha-\beta+\gamma-\frac{5}{2}} \times x^{1-2\alpha} z^{1-2\gamma} x_0^{1-2\alpha} z_0^{1-2\gamma} F_A^{(3)} \left(-\alpha + \beta - \gamma + \frac{5}{2}; 1 - \alpha, \beta, 1 - \gamma; 2 - 2\alpha, 2\beta, 2 - 2\gamma; \xi, \eta, \zeta \right), \quad (36)$$

$$q_7(x, y, z; x_0, y_0, z_0) = k_7 (r^2)^{-\alpha+\beta+\gamma-\frac{5}{2}} \times y^{1-2\beta} z^{1-2\gamma} y_0^{1-2\beta} z_0^{1-2\gamma} F_A^{(3)} \left(\alpha - \beta - \gamma + \frac{5}{2}; \alpha, 1 - \beta, 1 - \gamma; 2\alpha, 2 - 2\beta, 2 - 2\gamma; \xi, \eta, \zeta \right), \quad (37)$$

$$q_8(x, y, z; x_0, y_0, z_0) = k_8 (r^2)^{\alpha+\beta+\gamma-\frac{7}{2}} x^{1-2\alpha} y^{1-2\beta} z^{1-2\gamma} x_0^{1-2\alpha} y_0^{1-2\beta} z_0^{1-2\gamma} \times F_A^{(3)} \left(-\alpha - \beta - \gamma + \frac{7}{2}; 1 - \alpha, 1 - \beta, 1 - \gamma; 2 - 2\alpha, 2 - 2\beta, 2 - 2\gamma; \xi, \eta, \zeta \right). \quad (38)$$

Here k_i ($i = \overline{1, 8}$) are constants. They will be determined when we solve boundary-value problems.

Let us show that the found solutions (31) - (38) have a singularity.

We choose a solution $q_1(x, y, z; x_0, y_0, z_0)$. For this aim we use expansion for hypergeometric function of Lauricella (15). As a result solution (31) can be written as follows

$$\begin{aligned} q_1(x, y, z; x_0, y_0, z_0) &= k_1 (r^2)^{-\alpha-\beta-\gamma-\frac{1}{2}} \sum_{l, m, n=0}^{\infty} \frac{\left(\alpha + \beta + \gamma + \frac{1}{2} \right)_{l+m+n} (\alpha)_{l+m} (\beta)_{l+n} (\gamma)_{m+n}}{(2\alpha)_{l+m} (2\beta)_{l+n} (2\gamma)_{m+n} l! m! n!} \\ &\times \left(1 - \frac{r_1^2}{r^2} \right)^{l+m} \left(1 - \frac{r_2^2}{r^2} \right)^{l+n} \left(1 - \frac{r_3^2}{r^2} \right)^{m+n} \\ &\times {}_2F_1 \left(\alpha + \beta + \gamma + \frac{1}{2} + l + m, \alpha + l + m; 2\alpha + l + m; 1 - \frac{r_1^2}{r^2} \right) \\ &\times {}_2F_1 \left(\alpha + \beta + \gamma + \frac{1}{2} + l + m + n, \beta + l + n; 2\beta + l + n; 1 - \frac{r_2^2}{r^2} \right) \\ &\times {}_2F_1 \left(\alpha + \beta + \gamma + \frac{1}{2} + l + m + n, \gamma + m + n; 2\gamma + m + n; 1 - \frac{r_3^2}{r^2} \right). \end{aligned} \quad (39)$$

Using formula (5), we rewrite (39) as

$$q_1(x, y, z; x_0, y_0, z_0) = k_1 r^{-1} (r_1^2)^{-\alpha} (r_2^2)^{-\beta} (r_3^2)^{-\gamma} f(r^2, r_1^2, r_2^2, r_3^2), \quad (40)$$

where

$$\begin{aligned}
f(r^2, r_1^2, r_2^2, r_3^2) &= \sum_{l, m, n=0}^{\infty} \frac{\left(\alpha + \beta + \gamma + \frac{1}{2}\right)_{l+m+n} (\alpha)_{l+m} (\beta)_{l+n} (\gamma)_{m+n}}{(2\alpha)_{l+m} (2\beta)_{l+n} (2\gamma)_{m+n} l! m! n!} \\
&\times \left(\frac{r^2}{r_1^2} - 1\right)^{l+m} \left(\frac{r^2}{r_2^2} - 1\right)^{l+n} \left(\frac{r^2}{r_3^2} - 1\right)^{m+n} \\
&\times {}_2F_1\left(\alpha - \beta - \gamma - \frac{1}{2}, \alpha + l + m; 2\alpha + l + m; 1 - \frac{r^2}{r_1^2}\right) \\
&\times {}_2F_1\left(\beta - \alpha - \gamma - \frac{1}{2} - m, \beta + l + n; 2\beta + l + n; 1 - \frac{r^2}{r_2^2}\right) \\
&\times {}_2F_1\left(\gamma - \alpha - \beta - \frac{1}{2} - l, \gamma + m + n; 2\gamma + m + n; 1 - \frac{r^2}{r_3^2}\right).
\end{aligned} \tag{41}$$

Below we show that $f(r^2, r_1^2, r_2^2, r_3^2)$ will be constant at $r \rightarrow 0$.

For this aim we use an equality (4). Then we get

$$\begin{aligned}
&{}_2F_1\left(\alpha - \beta - \gamma - \frac{1}{2}, \alpha + l + m; 2\alpha + l + m; 1 - \frac{r^2}{r_1^2}\right)\Big|_{r=0} \\
&= \frac{\Gamma\left(\beta + \gamma + \frac{1}{2}\right) \Gamma(2\alpha + l + m)}{\Gamma(\alpha) \Gamma\left(\alpha + \beta + \gamma + \frac{1}{2} + l + m\right)}.
\end{aligned}$$

Considering first formula in (3), we obtain

$$\begin{aligned}
&{}_2F_1\left(\alpha - \beta - \gamma - \frac{1}{2}, \alpha + l + m; 2\alpha + l + m; 1 - \frac{r^2}{r_1^2}\right)\Big|_{r=0} \\
&= \frac{\Gamma\left(\beta + \gamma + \frac{1}{2}\right) \Gamma(2\alpha) (2\alpha)_{l+m}}{\Gamma(\alpha) \Gamma\left(\alpha + \beta + \gamma + \frac{1}{2}\right) \left(\alpha + \beta + \gamma + \frac{1}{2}\right)_{l+m}}.
\end{aligned} \tag{42}$$

Similarly we get

$$\begin{aligned}
&{}_2F_1\left(\beta - \alpha - \gamma - \frac{1}{2} - m, \beta + l + n; 2\beta + l + n; 1 - \frac{r^2}{r_2^2}\right)\Big|_{r=0} \\
&= \frac{\Gamma(2\beta) \Gamma\left(\alpha + \gamma + \frac{1}{2}\right) (2\beta)_{l+n} \left(\alpha + \gamma + \frac{1}{2}\right)_m}{\Gamma(\beta) \Gamma\left(\alpha + \beta + \gamma + \frac{1}{2}\right) \left(\alpha + \beta + \gamma + \frac{1}{2}\right)_{l+m+n}},
\end{aligned} \tag{43}$$

$$\begin{aligned}
&{}_2F_1\left(\gamma - \alpha - \beta - \frac{1}{2} - l, \gamma + m + n; 2\gamma + m + n; 1 - \frac{r^2}{r_3^2}\right)\Big|_{r=0} \\
&= \frac{\Gamma(2\gamma) \Gamma\left(\alpha + \beta + \frac{1}{2}\right) (2\gamma)_{m+n} \left(\alpha + \beta + \frac{1}{2}\right)_l}{\Gamma(\gamma) \Gamma\left(\alpha + \gamma + \beta + \frac{1}{2}\right) \left(\alpha + \gamma + \beta + \frac{1}{2}\right)_{l+m+n}}.
\end{aligned} \tag{44}$$

Taking (42) - (44) into account from (41) at $r \rightarrow 0$ we have

$$f(0, r_1^2, r_2^2, r_3^2) = \frac{\Gamma(2\alpha) \Gamma(2\beta) \Gamma(2\gamma) \Gamma\left(\alpha + \beta + \frac{1}{2}\right) \Gamma\left(\alpha + \gamma + \frac{1}{2}\right) \Gamma\left(\beta + \gamma + \frac{1}{2}\right)}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\gamma) \Gamma^3\left(\alpha + \beta + \gamma + \frac{1}{2}\right)} \cdot \sum_{l,m,n=0}^{\infty} \frac{\left(\alpha + \gamma + \frac{1}{2}\right)_m \left(\alpha + \beta + \frac{1}{2}\right)_l (\alpha)_{l+m} (\beta)_{l+n} (\gamma)_{m+n}}{\left(\alpha + \gamma + \beta + \frac{1}{2}\right)_{l+m+n} \left(\alpha + \beta + \gamma + \frac{1}{2}\right)_{l+m} l!m!n!}. \quad (45)$$

We use formulas in (3) to simplify (45). Since

$$\begin{aligned} & \sum_{l,m,n=0}^{\infty} \frac{\left(\alpha + \gamma + \frac{1}{2}\right)_m \left(\alpha + \beta + \frac{1}{2}\right)_l (\alpha)_{l+m} (\beta)_{l+n} (\gamma)_{m+n}}{\left(\alpha + \gamma + \beta + \frac{1}{2}\right)_{l+m+n} \left(\alpha + \beta + \gamma + \frac{1}{2}\right)_{l+m} l!m!n!} \\ &= \sum_{l,m,n=0}^{\infty} \frac{\left(\alpha + \gamma + \frac{1}{2}\right)_m \left(\alpha + \beta + \frac{1}{2}\right)_l (\alpha)_l (\alpha + l)_m (\beta)_l (\beta + l)_n (\gamma)_m (\gamma + m)_n}{\left(\alpha + \gamma + \beta + \frac{1}{2} + l + m\right)_n \left(\left(\alpha + \beta + \gamma + \frac{1}{2}\right)_{l+m}\right)^2 l!m!n!} \\ &= \sum_{l,m=0}^{\infty} \frac{\left(\alpha + \gamma + \frac{1}{2}\right)_m \left(\alpha + \beta + \frac{1}{2}\right)_l (\alpha)_l (\alpha + l)_m (\beta)_l (\gamma)_m}{\left(\left(\alpha + \beta + \gamma + \frac{1}{2}\right)_{l+m}\right)^2 l!m!} \cdot \sum_{l,m=0}^{\infty} \frac{(\beta + l)_n (\gamma + m)_n}{\left(\alpha + \gamma + \beta + \frac{1}{2} + l + m\right)_n n!} \\ &= \frac{\Gamma\left(\alpha + \beta + \gamma + \frac{1}{2}\right) \Gamma\left(\alpha + \frac{1}{2}\right)}{\Gamma\left(\alpha + \gamma + \frac{1}{2}\right) \Gamma\left(\alpha + \beta + \frac{1}{2}\right)} \sum_{l=0}^{\infty} \frac{(\alpha)_l (\beta)_l}{\left(\alpha + \beta + \gamma + \frac{1}{2}\right)_l l!} \sum_{m=0}^{\infty} \frac{(\gamma)_m (\alpha + l)_m}{\left(\alpha + \beta + \gamma + \frac{1}{2} + l\right)_m m!} \\ &= \frac{\Gamma^2\left(\alpha + \beta + \gamma + \frac{1}{2}\right) \Gamma\left(\alpha + \frac{1}{2}\right) \Gamma\left(\beta + \frac{1}{2}\right)}{\Gamma^2\left(\alpha + \beta + \frac{1}{2}\right) \Gamma\left(\alpha + \gamma + \frac{1}{2}\right) \Gamma\left(\beta + \gamma + \frac{1}{2}\right)} \sum_{l=0}^{\infty} \frac{(\alpha)_l (\beta)_l}{\left(\alpha + \beta + \gamma + \frac{1}{2}\right)_l l!} \\ &= \frac{\Gamma^2\left(\alpha + \gamma + \beta + \frac{1}{2}\right) \sqrt{\pi}}{\Gamma\left(\alpha + \beta + \frac{1}{2}\right) \Gamma\left(\alpha + \gamma + \frac{1}{2}\right) \Gamma\left(\beta + \gamma + \frac{1}{2}\right)}, \end{aligned}$$

from (45) follows

$$f(0, r_1^2, r_2^2, r_3^2) = \frac{\Gamma(2\alpha) \Gamma(2\beta) \Gamma(2\gamma) \sqrt{\pi}}{\Gamma(\alpha) \Gamma(\beta) \Gamma(\gamma) \Gamma\left(\alpha + \beta + \gamma + \frac{1}{2}\right)}. \quad (46)$$

Expressions (46) and (40) gives us possibility to conclude that the solution $q_1(x, y, z; x_0, y_0, z_0)$ reduce to infinity of the order r^{-1} at $r \rightarrow 0$. Similarly it is possible to be convinced that solutions $q_i(x, y, z; x_0, y_0, z_0)$, $i = 2, 3, \dots, 8$ also reduce to infinity of the order r^{-1} when $r \rightarrow 0$.

It is directly checkable that constructed functions (31) - (38) possess properties

$$\begin{aligned} x^{2\alpha} \frac{\partial}{\partial x} q_1|_{x=0} &= 0, \quad y^{2\beta} \frac{\partial}{\partial y} q_1|_{y=0} = 0, \quad z^{2\gamma} \frac{\partial}{\partial z} q_1|_{z=0} = 0, \\ q_2|_{x=0} &= 0, \quad y^{2\beta} \frac{\partial}{\partial y} q_2|_{y=0} = 0, \quad z^{2\gamma} \frac{\partial}{\partial z} q_2|_{z=0} = 0, \end{aligned}$$

$$\begin{aligned}
x^{2\alpha} \frac{\partial}{\partial x} q_3|_{x=0} &= 0, \quad q_3|_{y=0} = 0, \quad z^{2\gamma} \frac{\partial}{\partial z} q_3|_{z=0} = 0, \\
x^{2\alpha} \frac{\partial}{\partial x} q_4|_{x=0} &= 0, \quad y^{2\beta} \frac{\partial}{\partial y} q_4|_{y=0} = 0, \quad q_4|_{z=0} = 0, \\
q_5|_{x=0} &= 0, \quad q_5|_{y=0} = 0, \quad z^{2\gamma} \frac{\partial}{\partial z} q_5|_{z=0} = 0, \\
q_6|_{x=0} &= 0, \quad y^{2\beta} \frac{\partial}{\partial y} q_6|_{y=0} = 0, \quad q_6|_{z=0} = 0, \\
x^{2\alpha} \frac{\partial}{\partial x} q_7|_{x=0} &= 0, \quad q_7|_{y=0} = 0, \quad q_7|_{z=0} = 0, \\
q_8|_{x=0} &= 0, \quad q_8|_{y=0} = 0, \quad q_8|_{z=0} = 0.
\end{aligned}$$

Let be convinced that

$$x^{2\alpha} \frac{\partial}{\partial x} q_1|_{x=0} = 0. \quad (47)$$

Using formula of differentiation for Lauricella HF of three variables (16) we have

$$\begin{aligned}
\frac{\partial}{\partial x} q_1(x, y, z; x_0, y_0, z_0) &= -2(x - x_0) \left(\alpha + \beta + \gamma + \frac{1}{2} \right) k_1(r^2)^{-\alpha-\beta-\gamma-\frac{3}{2}} \\
&\times \left[F_A^{(3)} \left(\alpha + \beta + \gamma + \frac{3}{2}; \alpha, \beta, \gamma; 2\alpha, 2\beta, 2\gamma; \xi, \eta, \zeta \right) \right. \\
&+ \frac{\xi}{2} F_A^{(3)} \left(\alpha + \beta + \gamma + \frac{3}{2}; \alpha + 1, \beta, \gamma; 2\alpha + 1, 2\beta, 2\gamma; \xi, \eta, \zeta \right) \\
&+ \frac{\eta}{2} F_A^{(3)} \left(\alpha + \beta + \gamma + \frac{3}{2}; \alpha, \beta + 1, \gamma; 2\alpha, 2\beta + 1, 2\gamma; \xi, \eta, \zeta \right) \\
&\left. + \frac{\zeta}{2} F_A^{(3)} \left(\alpha + \beta + \gamma + \frac{3}{2}; \alpha, \beta, \gamma + 1; 2\alpha, 2\beta, 2\gamma + 1; \xi, \eta, \zeta \right) \right] \\
&- 4x_0 \left(\alpha + \beta + \gamma + \frac{1}{2} \right) k_1(r^2)^{-\alpha-\beta-\gamma-\frac{3}{2}} F_A^{(3)} \left(\alpha + \beta + \gamma + \frac{3}{2}; \alpha, \beta, \gamma; 2\alpha, 2\beta, 2\gamma; \xi, \eta, \zeta \right).
\end{aligned}$$

From the last equality one can easily be convinced that (47) is true.

Remark. *Afore-mentioned properties of found fundamental solutions will be used at solving various boundary-value problems for the equation (1). For instance, fundamental solution (31) with its properties will be used at solving problem N for the equation (1).*

Acknowledgement.

The authors are grateful to Professor R.Gilbert for his kind interest and given materials on a theme of the paper.

References

- [1] A. Altin, Some expansion formulas for a class of singular partial differential equations. *Proc. Amer. Math. Soc.* Volume 85, Number 1 (1982), 42-46.
- [2] P. Appell, J. Kampe de Fériet, *Fonctions Hypergeometriques et Hyperspheriques; Polynomes d'Hermite*, Gauthier - Villars. Paris, 1926.
- [3] J. Barros-Neto and I. M. Gelfand, Fundamental solutions for the Tricomi operator, *Duke Math. J.* Volume 98, Number 3 (1999), 465-483.
- [4] J. Barros-Neto and I. M. Gelfand. Fundamental solutions for the Tricomi operator II, *Duke Math. J.* Volume 111, Number 3 (2002), 561-584.

- [5] J. Barros-Neto and I. M. Gelfand, Fundamental solutions for the Tricomi operator III, *Duke Math.J.* Volume 128, Number 1 (2005), 119-140.
- [6] Erdelyi, A. and Bateman G.: *Higher transcendental functions*, vol. 1. Moscow: Nauka. 1973. (Russian translation)
- [7] A. J. Fryant, Growth and complete sequences of generalized bi-axially symmetric potentials. *J. Differential Equations*, Volume 31, Number 2 (1979), 155-164.
- [8] M.A. Golberg and C.S. Chen, The method of fundamental solutions for potential, Helmholtz and diffusion problems. In Boundary Integral Methods - Numerical and Mathematical Aspects, (Ed. by M.A. Golberg), pp. 103-176, *Comput. Mech. Publ.*, 1998.
- [9] A. Hasanov and H. M. Srivastava, Some decomposition formulas associated with the Lauricella function $F_A^{(r)}$ and other multiple hypergeometric functions, *Appl. Math. Lett.* Volume 19, Number 2 (2006), 113-121.
- [10] A. Hasanov and H. M. Srivastava, Decomposition formulas associated with the Lauricella multivariable hypergeometric functions, *Computers and mathematics with applications*, Volume 53, (2007), 1119-1228.
- [11] M. Itagaki, Higher order three-dimensional fundamental solutions to the Helmholtz and the modified Helmholtz equations, *Engineering Analysis with Boundary elements*. Volume 15 (1995), 289-293.
- [12] J. Leray, Un prolongement de la transformation de Laplace qui transforme la solution unitaires d'un opérateur hyperbolique en sa solution élémentaire (Problème de Cauchy, IV), *Bull. Soc. Math. France*. Volume 90 (1962), 39-156.
- [13] P. A. McCoy, Polynomial approximation and growth of generalized axisymmetric potentials, *Canad. J. Math.* Volume 31, Number 1 (1979), 49-59.
- [14] P. Kumar, Approximation of growth numbers generalized bi-axially symmetric potentials, *Fasc. Math.* Volume 35 (2005), 51-60.
- [15] N. Radzhabov, Integral representations and boundary value problems for an equation of Helmholtz type with several singular surfaces, *Analytic methods in the theory of elliptic equations*, pp. 34-46, "Nauka", *Sibirsk. Otdel. Novosibirsk*, 1982.
- [16] R. J. Weinacht, Fundamental solutions for a class of singular equations, *Contrib. Differential Equations*, Volume 3 (1964), 43-55.
- [17] A. Weinstein, Discontinuous integrals and generalized potential theory, *Trans. Amer. Math. Soc.*, Volume 63 (1946), 342-354.
- [18] A. Weinstein, Generalized axially symmetric potential theory, *Bull. Amer. Math. Soc.*, Volume 59 (1953), 20-38.